

Please type a plus (+) inside this box → ☐

PTO/SB/29 (12/97)

Approved for use through 09/30/00. OMB 0651-0032
Patent and Trademark Office: U.S. DEPARTMENT OF COMMERCE

Under the Paperwork Reduction Act of 1995, no persons are required to respond to a collection of information unless it displays a valid OMB control number.

**UTILITY
PATENT APPLICATION
TRANSMITTAL**

(Only for new nonprovisional applications under 37 CFR 1.53(b))

Attorney Docket No. 8824-016

Total Pages 20

First Named Inventor or Application Identifier

Amy Arrow

Express Mail Label No.

EM 061 017 733 US

APPLICATION ELEMENTS

See MPEP chapter 600 concerning utility patent application contents.

ADDRESS TO:

Assistant Commissioner for Patents
Box Patent Application
Washington, DC 20231

1. ☒ Fee Transmittal Form
Submit an original, and a duplicate for fee processing
2. ☒ Specification [Total Pages 20]
(preferred arrangement set forth below)
- Descriptive title of the Invention
 - Cross Reference to Related Applications
 - Statement Regarding Fed sponsored R&D
 - Reference to Microfiche Appendix
 - Background of the Invention
 - Brief Summary of the Invention
 - Brief Description of the Drawings *(if filed)*
 - Detailed Description of the Invention *(including drawings, if filed)*
 - Claim(s)
 - Abstract of the Disclosure

3. ☐ Drawing(s) (35 USC 113) (Figs. 1-18) [Total Sheets 0]

4. ☒ Oath or Declaration (Unexecuted) [Total Sheets 2]

a. ☐ Newly executed (original or copy)

b. ☐ Copy from a prior application (37 CFR 1.63(d))
(for continuation/divisional with Box 17 completed)
[Note Box 5 below]

i. ☐ **DELETION OF INVENTOR(S)**

Signed statement attached deleting inventor(s) named in the prior application, see 37 CFR 1.63(d)(2) and 1.33 (b).

5. ☐ Incorporation By Reference *(useable if Box 4b is checked)*
The entire disclosure of the prior application, from which a copy of the oath or declaration is supplied under Box 4b, is considered as being part of the disclosure of the accompanying application and is hereby incorporated by reference therein.

6. ☐ Microfiche Computer Program *(Appendix)*
7. ☐ Nucleotide and/or Amino Acid Sequence Submission
(if applicable, all necessary)
- a. ☐ Computer Readable Copy
 - b. ☐ Paper Copy (identical to computer copy)
 - c. ☐ Statement verifying identity of above copies

ACCOMPANYING APPLICATION PARTS

8. ☐ Assignment Papers (cover sheet & document(s))
9. ☐ 37 CFR 3.73(b) Statement ☐ Power of Attorney
(when there is an assignee)
10. ☐ English Translation Document *(if applicable)*
11. ☐ Information Disclosure Statement (IDS)/PTO-1449 ☐ Copies of IDS Citations
12. ☐ Preliminary Amendment
13. ☒ Return Receipt Postcard (MPEP 503)
(Should be specifically itemized)
14. ☐ Small Entity ☐ Statement filed in prior application, Statement(s) Status still proper and desired
15. ☐ Certified Copy of Priority Document(s)
(if foreign priority is claimed)
16. ☐ Other:

17. If a CONTINUING APPLICATION, check appropriate box and supply the requisite information:

☒ Continuation ☐ Divisional ☐ Continuation-in-part (CIP) of prior application No: 08/754,580 filed 11/21/1996

18. CORRESPONDENCE ADDRESS

☒ Customer Number or Bar Code Label

20583
(Insert Customer No. or Attach bar code label here)

or ☐ Correspondence address below

NAME

Pennie & Edmonds LLP

ADDRESS

1155 Avenue of the Americas

CITY

New York

STATE

NY

ZIP CODE

10036-2711

COUNTRY

USA

TELEPHONE

212 790-9090

FAX

212 869-9741

Burden Hour Statement: This form is estimated to take 0.2 hours to complete. Time will vary depending upon the needs of the individual case. Any comments on the amount of time you are required to complete this form should be sent to the Chief Information Officer, Patent and Trademark Office, Washington, DC 20231. DO NOT SEND FEES OR COMPLETED FORMS TO THIS ADDRESS. SEND TO: Assistant Commissioner for Patents, Box Patent Application, Washington, DC 20231.

IN THE UNITED STATES PATENT AND TRADEMARK OFFICEPrior application: Examiner J. LeguyaderArt Unit 1635

Assistant Commissioner for Patents
 Box PATENT APPLICATION
 Washington, D.C. 20231

Sir:

This is a request for filing a ☒ continuation ☐ divisional application under 37 CFR § 1.53(b), of pending prior application no. 08/754,580 filed on November 21, 1996

of Amy Arrow, Roderic M.K. Dale; Tod Mitchell Woolf
 (inventor(s) currently of record in prior application)

for THREE COMPONENT CHIMERIC ANTISENSE OLIGONUCLEOTIDES
 (title of invention)

1. ☒ The filing fee is calculated below:

PATENT APPLICATION FEE VALUE


TYPE	NO. FILED	LESS	EXTRA	EXTRA RATE	FEE
Total Claims	19	-20	0	\$18.00 each	0.00
Independent	3	-3	0	\$76.00 each	0.00
Basic Fee					760.00
Multiple Dependency Fee If Applicable (\$270.00)					
Total					760.00
50% Reduction for Independent Inventor, Nonprofit Organization or Small Business Concern					-
Total Filing Fee					\$ 760.00

2. ☒ Please charge the required fee to Pennie & Edmonds LLP Deposit Account No. 16-1150. A copy of this sheet is enclosed.
3. ☒ Amend the specification by inserting before the first line the following sentence: This is a ☒ continuation, ☐ division, of application no. 08/754,580, filed November 21, 1996.
- 4a. ☐ Transfer the drawing from the prior application to this application.

- 4b. ☐ New formal drawings are enclosed.
- 4c. ☐ Informal drawings are enclosed.
- 5a. ☐ Priority of application no. filed on is claimed under 35 U.S.C. §119.
- 5b. ☐ The certified copy has been filed in prior application no.
- 6. ☐ The prior application is assigned of record to
- 7a. ☒ The Power of Attorney appears in the original papers in the prior application no. 08/754,580, filed November 21, 1996.
- 7b. ☐ Since the Power of Attorney does not appear in the original papers, a copy of the Power in prior application no. $\phi\phi$, filed $\phi\phi$ is enclosed.
- 8. ☐ This application contains nucleic acid and/or amino acid sequences required to be disclosed in a Sequence Listing under 37 CFR §§1.821-1.825. It is requested that the Sequence Listing in computer readable form from prior application no. $\phi\phi$, filed on $\phi\phi$ be made a part of the present application as provided for by 37 C.F.R. §1.821(e). The sequences disclosed therein are the same as the sequences disclosed in this application. A copy of the paper Sequence Listing from application no. $\phi\phi$ is enclosed.
- 9. ☐ The undersigned states, under 37 C.F.R. §1.821(f), that the content of the enclosed paper Sequence Listing from application no. $\phi\phi$ is the same as the content of the computer readable form submitted in application no. $\phi\phi$.
- 10. ☒ Additional enclosures or instructions: Utility Patent Application Transmittal; a copy of the specification.

Respectfully submitted,

December 15, 1998
(date)


Thomas E. Friebe (signature) 29,258
PENNIE & EDMONDS LLP (Reg No.)
1155 Avenue of the Americas
New York, NY 10036
Tel. No. (212) 790-9090

THREE COMPONENT CHIMERIC ANTISENSE OLIGONUCLEOTIDES

1. FIELD OF THE INVENTION

This invention relates to antisense oligonucleotides that target mRNAs in cells as substrates for the cellular enzyme RNase H and thereby cause specific degradation of the targeted mRNA. The oligonucleotides have three components: a RNase H activating region, a complementarity region and 3' and 5' ends. The invention optimizes each of the components to resist intracellular nucleases, to increase hybridization to target mRNA, to specifically inactivate target mRNA in cells, and to decrease cytotoxicity.

2. BACKGROUND TO THE INVENTION

Antisense polynucleotides are useful for specifically inhibiting unwanted gene expression in mammalian cells. They can be used to hybridize to and inhibit the function of an RNA, typically a messenger RNA, by activating RNase H.

The use of antisense oligonucleotides has emerged as a powerful new approach for the treatment of certain diseases. The preponderance of the work to date has focused on the use of antisense oligonucleotides as antiviral agents or as anticancer agents (Wickstrom, E., Ed., Prospects for Antisense Nucleic Acid Therapy of Cancer and AIDS, New York: Wiley-Liss, 1991; Crooke, S.T. and Lebleu, B., Eds., Antisense Research and Applications, Boca Raton: CRC Press, 1993, pp. 154-182; Baserga, R. and Denhardt, D.T., 1992, Antisense Strategies, New York: The New York Academy of Sciences, Vol. 660; Murray, J.A.H., Ed., Antisense RNA and DNA, New York: Wiley-Liss, 1993).

There have been numerous disclosures of the use of antisense oligonucleotides as antiviral agents. For example, Agrawal et al. report phosphoramidate and phosphorothioate oligonucleotides as antisense inhibitors of HIV. Agrawal et al., Proc. Natl. Acad. Sci. USA 85, 7079-7083 (1988). Zamecnik et al. disclose antisense oligonucleotides as

inhibitors of Rous sarcoma virus replication in chicken fibroblasts. Zamecnik et al., Proc. Natl. Acad. Sci. USA 83, 4143-4146 (1986).

The principal mechanism by which antisense oligonucleotides affect the level of the target RNA is by activation of RNase H, which cleaves the RNA strand of DNA/RNA hybrids. Both phosphodiester and phosphorothioate-linked DNA activates endogenous RNase H, thereby cleaving the targeted RNA (Agrawal, S., et al., Proc. Natl. Acad. Sci. USA 87, 1101-5 (1990); Woolf, T.M., et al., Nucleic Acids Res. 18, 1763-9 (1990)). However, phosphodiester-linked DNA is rapidly degraded by cellular nucleases and, with the exception of the phosphorothioate-linked DNA, nuclease resistant, non-naturally occurring DNA derivatives do not activate RNase H when hybridized to RNA. While phosphorothioate DNA has the advantage of activating RNase H, phosphorothioate-linked DNA has non-specific cytotoxic effects and also has reduced affinity for RNA (Stein, C.A., et al., Aids Res Hum Retroviruses 5, 639-46 (1989); Woolf, T.M., et al., Nucleic Acids Res. 18, 1763-9 (1990); Kawasaki, A.M., et al., J. Med. Chem. 36, 831-41 (1993)).

Chimeric antisense oligos that have a short stretch of phosphorothioate DNA (3-9 bases) have been used to obtain RNase H-mediated cleavage of the target RNA (Dagle, J.M., et al., Nucleic Acids Res. 18, 4751-7 (1990); Agrawal, S., et al., Proc. Natl. Acad. Sci. USA 87, 1401-5 (1990); Monia, B.P. et al., 1993, J. Biol. Chem. 268, 14514). A minimum of 3 DNA bases is required for activation of bacterial RNase H (Futdon, P.J., et al., Nucleic Acids Res. 17, 9193-9204; Quartin, R.S., et al., Nucleic Acids Res. 17, 7235-7262) and a minimum of 5 bases is required for mammalian bacterial RNase H activation (Monia, B.P., et al., J. Biol. Chem. 268, 14514-14522 (1993)). In these chimeric oligonucleotides there is a central region that forms a substrate for RNase H that is flanked by hybridizing "arms," comprised of modified nucleotides that do not form substrates for RNase H. Alternatively, extracellular tests using a RNase H-containing

HeLa cell extract have been reported wherein the RNase H activating region was placed on the 5' or 3' side of the oligomer. Specifically these tests reported that a 5' or 3' terminal RNase H activating region composed of phosphodiester 2'-deoxynucleotides joined to a methylphosphonate-linked complementarity region was fully active, but that a 5' terminal RNase H-activating region composed of phosphorothioate 2'-deoxynucleotides joined to a methylphosphonate-linked complementarity region was only partially active. See Col 10, U.S. Pat. No. 5,220,007 to T. Pederson et al.

- 2'-O-Methyl or 2'-fluoro modified nucleotides have been used for the hybridizing arms of chimeric oligos. Inoue, H., et al., 1987, Nucleic Acids Res. 15, 6131-48. The 2'-O-Methyl group increases the affinity of the oligomer for the targeted RNA and increases the activity of the oligomer in cell culture. However, 2'-O-Methyl bases with phosphodiester linkages are degraded by exonucleases and so are not suitable for use in cell or therapeutic applications of antisense.
- Shibahara, S., et al., 1989, Nucleic Acids Res. 17, 239-52. Phosphorothioate 2'-O-Methyl nucleotides are resistant to nucleases as shown in the uniformly phosphorothioate modified oligos described by Monia B.P., et al., 1993, J. Biol. Chem. 268, 14514-14522 and terminal phosphorothioate substituted, 2'-O-Methylribo-oligonucleotides, Shibahara, S., et al., 1989, Nucleic Acid Res. 17, 239-252. However, fully phosphorothioate substituted oligomers may cause non-specific effects including cell toxicity. Stein, C.A., et al., 1989, Aids Res. Hum. Retrov. 5, 639-646; Woolf, T.M., et al., 1990, Nucleic Acids Res. 18, 1763-69; Wagner, R.W., 1995, Antisense Res. Dev. 5, 113-115; Krieg, A.M., & Stein, C.A., 1995, Antisense Res. Dev. 5, 241. The effects of 2'-Fluoro-oligonucleotides on bacterial RNase H are discussed in Crooke, S.T. et al., 1995, Bioch. J. 312, 599-608 and Iwai, S. et al., 1995, FEBS Lett (Neth.) 368, 315-20.

Several other chemistries have been used to make the "arms" or regions of a chimeric oligomer that are not

substrates for RNase H. The first chimeric oligomers used methylphosphonate or phosphoramidate linkages in the arms (Dagle, J.M., Walder, J.A. & Weeks, K.L., Nucleic Acids Res. 18, 1751-7 (1990); Agrawal, S., et al., Proc. Natl. Acad.

5 Sci. USA 87, 1401-5 (1990). While these compounds functioned well in buffer systems and *Xenopus* oocytes, the arms decreased the hybrid affinity. This decrease in affinity dramatically reduces the activity of oligomers in mammalian cell culture.

10 A number of studies have been reported for the synthesis of ethylated and methylated phosphotriester oligonucleotides and their physico-chemical and biochemical evaluation.

Dinucleotides with methyl and ethyl triesters were shown to possess greater affinity towards polynucleotides possessing

15 complementary sequences (Miller, P.S., et al., J. Am. Chem. Soc. 93, 6657, (1971)). However, a few years ago, another group reported lack of, or poor binding affinity of heptethyl ester of oligothymidine with complementary polynucleotides (Pless, R.C., and Ts'O, P.O.P., Biochemistry 16, 1239-1250

20 (1977)). Phosphate methylated (P-methoxy) oligonucleotides were synthesized and found to possess resistance towards endonuclease digestion (Gallo, K.L., et al. Nucl. Acid Res. 18, 7405 (1986)). A P-methoxy 18-mer oligonucleotide was shown to have high T_m value in duplexes with natural DNA and

25 blocked to the DNA replication process at room temperature (Moody, H.M., et al., Nucl. Acid Res. 17, 4769-4782 (1989)).

Moody et al. stated that phosphate ethylated (P-ethoxy) oligonucleotides would have poor antisense properties. P-methoxy dimers of DNA bases were synthesized using transient

30 protecting group of FMOC for the exocyclic amino groups (Koole, L.H., et al., J. Org. Chem. 54, 1657-1664 (1989)).

Synthesis and physico-chemical properties of partial P-methoxy oligodeoxyribonucleotides were determined. Only the thymidine and cytidine oligomers with methyl phosphotriester

35 could be prepared satisfactorily due to difficulty in maintaining methyl triester intact. Furthermore, the methyl group was found to have destabilizing effect on the

hybridization properties of the modified oligomers with its complementary sequence by comparison with unmodified parent oligodeoxynucleotide (Vinogradov, S., Asseline, U., Thoung, N.T., Tet. Let. 34, 5899-5902 (1993)).

- 5 Other reports have suggested that P-methoxy oligonucleotides are preferable to P-ethoxy as antisense oligonucleotides because of p-methoxy oligonucleotides showed stronger hybridization than methyl phosphonate or P-ethoxy oligonucleotides (van Genderen, M.H.P., et al., Kon. Ned. Akad. van Wetensch. B90, 155-159 (1987); van Genderen, M.H.P., et al., Trav. Chim. Pays Bas 108, 28-35 (1989)). P-ethoxy oligonucleotides were reported by van Genderen et al. to hybridize poorly to DNA, for which reason they were regarded unfavorably as antisense oligonucleotides (Moody, 15 H.M., et al., Nucl. Acid Res. 17, 4769-4782 (1989)).

P-isopropoxyphosphoramidites have been synthesized from several nucleosides (Stec, W.J., et al., Tet. Let. 26, 2191-2194 (1985)), and a few short oligonucleotides containing P-isopropoxy phosphotriesters were synthesized, and

- 20 hybridization studies were carried out.

United States Patent No. 5,525,719 to Srivastava, S., and Raza, S.K., June 11, 1996, suggests antisense oligonucleotides consisting of 2'-O-Methyl nucleotides linked by phosphodiester and/or P-ethoxy or P-methoxy,

- 25 phosphotriester moieties.

Thus, currently there are no nucleic acid chemistries nor any chimeras that have been developed that optimally achieve all the features that are needed to provide an effective antisense oligonucleotide i.e. low toxicity, high 30 specificity, nuclease resistance, ease of synthesis, RNase H compatibility.

3. SUMMARY OF THE INVENTION

- The invention provides a class of oligonucleotide that 35 is optimized to target a specific RNA for RNase H degradation and to be itself resistant to degradation within in plasma and within eukaryotic, especially mammalian cells. The

oligonucleotides of the invention contain no naturally occurring 5'→3'-linked nucleotides. Rather, the invention provides oligonucleotides having two types of nucleotides: 2'-deoxyphosphorothioate, which activate RNase H, and 2'-modified nucleotides, which do not. The linkages between the 2'-modified nucleotides can be phosphodiester, phosphorothioate or P-ethoxyphosphodiester. Activation of RNase H is accomplished by a contiguous, RNase H-activating region, which contains between three and five 2'-deoxyphosphorothioate nucleotides to activate bacterial RNase H and between five and ten 2'-deoxyphosphorothioate nucleotides to activate eukaryotic and, particularly, mammalian RNase H. Protection from degradation is accomplished by making the 5' and 3' terminal bases highly nuclease resistant and, optionally, by placing a 3' terminal blocking group.

In a preferred embodiment the RNase H activating region, which is composed of highly nuclease resistant phosphorothioate nucleotides is placed at the 5' end of the oligonucleotide.

4. DETAILED DESCRIPTION OF THE INVENTION

4.1 THE STRUCTURE OF THE OLIGONUCLEOTIDES

An oligonucleotide of the invention is comprised of a 3'-terminal 5'→3'-linked nucleoside and from 11 to 59 5'→3' linked nucleotides, which nucleotides can be 2'-deoxynucleotides or 2'-modified nucleotides, modified to enhance the hybridization of the oligonucleotide to the target mRNA, such as 2'-fluoro, 2'-methoxy, 2'-ethoxy, 2'-methoxyethoxy, 2'-allyloxy ($-OCH_2CH=CH_2$) nucleotides (hereinafter "2'-modified nucleotides"). The 3' terminal nucleoside can, optionally, be 2'-modified nucleoside. Those skilled in the art appreciate that the 3'-OH of the 3' terminal base can but need not be esterified to a phosphate or phosphate analog. The 3' terminal residue is referred to as a nucleoside even though it may be a nucleotide.

50
45
40
35
30
25
20
15
10
5
The internucleotide linkages of an oligonucleotide of the invention can be phosphodiester, phosphorothioate or P-ethoxyphosphodiester moieties. The oligonucleotide has a 3' terminus and a 5' terminus that are protected from nuclease attack. The 3' terminus is protected by having the 3' most 5'→3' linkage or linkages be a phosphorothioate or a P-alkyloxyphosphotriester linkage and/or by having a substituted 3' terminal hydroxyl, e.g., a 3'→3' linked nucleotide, wherein the alkyloxy radical is methoxy, ethoxy or isopropoxy and, preferably, ethoxy. Preferably two or three 3' terminal internucleotide linkages are phosphorothioate or a P-alkyloxyphosphotriester linkages. To reduce nuclease degradation, the 5' most 3'→5' linkage preferably should be a phosphorothioate linkage or P-alkyloxyphosphotriester linkage. Preferably, the two 5' most 3'→5' linkages should be phosphorothioate linkages or P-ethoxyphosphotriester linkages. Optionally, the 5'-terminal hydroxyl moiety can be esterified with a phosphorus containing moiety, e.g., phosphate, phosphorothioate or P-ethoxyphosphate, without limitation.

The 3' terminal 5'→3'-linked nucleoside has a 3'-OH that can be optionally substituted by a blocking moiety that prevents 3'-exonuclease degradation of the oligonucleotide. In one embodiment, the 3'-hydroxyl is esterified to a nucleotide through a 3'→3' internucleotide linkage. Optionally, the 3'→3' linked nucleotide at the 3' terminus can be linked by a phosphorothioate moiety.

In a preferred embodiment, the oligonucleotide contains, exclusive of an optional blocking nucleotide, between 15 and 50 bases and more preferably between 20 and 30 bases and in a most preferred embodiment the oligonucleotide is 25 bases in length. The oligonucleotide of the invention contains a single contiguous RNase H-activating region of between three to ten 2'-deoxyphosphorothioate nucleotides. The length of the RNase H activating region to activate bacterial RNase H is preferably between three and five nucleotides; to activate a eukaryotic RNase H it is between five and ten nucleotides.

The preferred length of the RNase H-activating region for the activation of mammalian RNase H is nine nucleotides.

All 5'→3' linked nucleotides of the oligonucleotide that are not a part of the RNase H-activating region are 2'-

5 modified nucleotides, which contribute to the target binding and form the complementarity determining region. The complementarity region can be a contiguous region or can be divided by the RNase H-activating region. In the preferred embodiment the complementarity region is a contiguous region, 10 and more preferably is 3' to the RNase H-activating region.

In a preferred embodiment all bases except the from one to three 3' most nucleotides and nucleoside, the 5' terminal nucleotide and RNase H activating region nucleotides are phosphodiester linked. Large amounts of contiguous 15 phosphorothioate linkages are detrimental to the function of the oligonucleotides of the invention. Preferably, therefore, the oligonucleotides contain not more than ten contiguous phosphorothioate linkages.

20 4.2 THE SYNTHESIS OF THE OLIGONUCLEOTIDES

The oligonucleotides of the invention can be synthesized by solid phase or liquid phase nucleotide synthesis, however, synthesis by solid phase techniques is preferred.

Phosphodiester and phosphorothioate linked oligonucleotides 25 can be synthesized, using standard reagents and protocols, on an automated synthesizer utilizing methods that are well known in the art, such as, for example, those disclosed in Stec et al., J. Am. Chem. Soc. 106, 6077-6089 (1984); Stec et al., J. Org. Chem. 50(20), 3908-3913 (1985); Stec et al., J. Chromatog. 326, 263-280 (1985); LaPlanche et al., Nuc. Acid. Res. 14, 9081-9093 (1986); and Fasman, G.D., Practical Handbook of Biochemistry and Molecular Biology 1989, CRC Press, Boca Raton, Florida, herein incorporated by reference.

The synthesis of 2'-O-alkyl-oligoribonucleotides, where 35 the alkyl groups are methyl, butyl, allyl or 3,3-dimethylallyl is reviewed by Lamond, Biochem. Soc. Trans. 21, 1-8 (1993). Intermediates that are useful in the synthesis

of 2'-O-methyl oligoribonucleotides are described in U.S. Patents No. 5,013,830, No. 5,525,719 and No. 5,214,135, which are hereby incorporated by reference.

The synthesis of 2'-fluorophosphodiester and 2'-
5 fluorophosphorothioate oligonucleotides can be performed according to teaching of Kawasaki, A.M., et al., 1993, J. Med. Chem. 36, 831-41 and WO 92/03568; the synthesis of P-alkyloxyphosphotriester-linked oligonucleotides and 2'-modified oligonucleotides can be performed according to U.S.
10 Patent No. 5,525,719, each of which is incorporated herein by reference. The synthesis of phosphorothioate oligodeoxynucleotides is taught by U.S. Patent No. 5,276,019 and No. 5,264,423, which is hereby incorporated by reference. Synthesis of 2'-substituted oligonucleotides can be performed
15 by variations on the techniques disclosed therein.

The synthesis of the oligonucleotides of the invention must be conducted with great attention to quality control. It is particularly important that the phosphorothioate linkages not be contaminated with phosphodiester linkages.
20 It is advisable to pre-test the individual reagent lots to ascertain that high coupling efficiency can be obtained therewith and to exercise all possible precautions to maintain anhydrous conditions.

The quality of the synthesis of oligonucleotides can be
25 verified by testing the oligonucleotides by capillary electrophoresis and denaturing strong anion HPLC (SAX-HPLC). The method of Bergot & Egan, 1992, J. Chrom. 599, 35-42 is suitable. SAX-HPLC is particularly useful to verify that the phosphorothioate nucleotides are completely thiolated, i.e.,
30 are not contaminated by a small percentage of phosphodiesters.

The synthesis of oligonucleotides having both phosphodiester and phosphorothioate linkages is associated with a side reaction whereby the phosphorothioate linkages
35 are oxidized by the standard I₂ treatments that are used to oxidize the cyanoethyl phosphoramidite. This problem can be minimized but not eliminated by reducing the concentration or

I₂ to as low as 0.001 M. Therefore, in a preferred embodiment, all phosphorothioates of the oligonucleotides of the invention are found at the 5'-end, so that no phosphorothioate bond is exposed to I₂.

5

4.3 THE USES OF THE OLIGONUCLEOTIDES

The oligonucleotides of the invention can be used as antisense oligonucleotides in a variety of *in vitro* experimental situations to specifically degrade an mRNA of
10 unknown function and thereby determine its physiologic function.

The oligonucleotides of the invention can be also used in clinical practice for any disease and against any target RNA for which antisense therapy is now known to be suitable
15 or which is yet to be identified. Medical conditions for which antisense therapy is reported to be suitable includes Respiratory Syncytial Virus infection, WO 95/22553 by Kilkuskie, Influenza Virus infection, WO 94/23028, and malignancies, WO 94/08003. Further examples of clinical uses
20 of antisense oligonucleotides are reviewed, in summary form, in Glaser, V., 1996, Genetic Engineering News 16, 1. Targets of antisense oligonucleotides under that are the subjects of clinical trials include protein kinase C α , ICAM-1, c-raf kinase, p53, c-myb and the bcr/abl fusion gene found in
25 chronic myelogenous leukemia.

5. EXAMPLES

5.1 EXPERIMENTAL CONDITIONS

The oligonucleotides of the invention are demonstrated
30 by a test transient expression system which includes an mRNA encoding the luciferase protein that has been modified to include a test sequence derived from the ras gene. The specific antisense effects of an oligonucleotide can be measured by comparing the luciferase production of the test
35 cells with the production of control cells having the same expression plasmid except for the absence of the ras-derived

sequence. The oligonucleotides of the invention tested have the sequence:

5'-TTGCCACACCGACGGCGCCACCA-3' (SEQ ID NO: 1)

The details of the assay are as follows:

- 5 Plasmid Constructs.** The plasmid used for the studies contained a portion of the ras gene sequence fused to luciferase (Monia, B.P., et al. J. Biol. Chem. 267, 19954-19962 (1992)). The control luciferase plasmids did not contain the ras target sequence.
- 10 Cell Culture Assay.** HeLa cells were grown to 40-90% confluence in DMEM/10% FBS, Supplemented with glutamine, penicillin and streptomycin on gelatin coated 24 well plates. The gelatin coating was necessary for cell to remain adherent during the transfections. Prior to transfection the cells
- 15** were washed twice with PBS (containing magnesium and calcium). LIPOFECTIN™ was mixed gently and 6.6μl was added for each milliliter of reduced serum medium (OPTI-MEM™, Gibco/BRL, Gaithersburg, MD). Oligomers were added from 50-100μM concentrated stock to make a master mixture. The Opti-
- 20** MEM/LIPOFECTIN/oligomer solution was added to the cells and incubated for 4 hours (≈0.5mls for one well of a 24 well plate).

A target transfection mixture was prepared by first diluting 5μl of lipofectin per ml of OPTI-MEM and mixing.

- 25** Next 5μg of luciferase target and 5μg of CMV β-galactosidase were added per milliliter of OPTI-MEM/LIPOFECTIN™ mixture. The transfection mixture was mixed gently and allowed to complex for about 15 minutes. The master mixture reduced error by assuring that the control and experimental cells
- 30** received the exact same cationic lipid/plasmid complex. The concentration of oligonucleotide in the culture medium was between 200nM and 400nM in all experiments. The oligonucleotide containing media was removed from the cells and replaced with growth media and incubated for an
- 35** additional 9-18 hours. The cell were rinsed with calcium and magnesium free media and the media was removed. The plates

were frozen at -70 for >20 minutes and 100-300 μ l of reporter lysis buffer (Promega, Madison WI) was added.

The cells were put through 2 more freeze thaw cycles, to assure complete lysis. Luciferase assays were preformed according to the manufacture's instructions (Promega, Madison WI) and luminescence was detected with a 96 well luminometer (Packard, Meriden CT). β -galactosidase assays were preformed (Galacton Plus, Tropix) according to manufactures instructions and detected on the Packard luminometer.

10

5.2 EXPERIMENTAL RESULTS

The results of luciferase assay are presented in Table I below. The results are reported as the percent specific inhibition which were calculated as

15 $100 \times (1 - (LUC_T/LUC_C)^{OLIGO} / (LUC_T/LUC_C)^{NO\ OLIGO})$; wherein LUC_T and LUC_C are the luciferase levels found in the cells transfected with luciferase plasmids containing and lacking the ras gene insert (SEQ ID NO: 1); and the superscripts "Oligo" and "No Oligo" refer to the presence and absence of antisense

20 oligonucleotides.

TABLE 1

Oligo	Formula	Specific inhibition
Controls ("C")		
25 C1	25Mo	26%
C2	25Ms	15%
C3	9Ds16Mo	15%
C4	9Do16MoInvT	0%
30 C5	9Dp16MoInvT	18%
C6	9Dp13Mo3Ms	14%
Controls with all "S"		
35 S1	25Ds	93%
S2	16Ms8DsD	100%
S3	8Ms9Ds7MsM	97%

S4	9Ds15MsM	95%
----	----------	-----

9Ds at 3' end ("3'I")

5	3'I1	InvTMs15Mo9DsInvT	59%
	3'I2	2Ms14Mo9DsInvT	57%
	3'I3	4Ms12Mo9DsInvT	65%

9Ds in Middle ("MI")

10	MI1	5Ms3Mo9Ds4Mo3MsM	64%
	MI2	2Ms6Mo9Ds7(MsMo)InvT	71%
	MI3	3Ms6Mo9Ds6MoMsInvT	87%

9Ds at 5'end ("5'I")

15	5'I1	9Ds16MoInvT	83%
	5'I2	9Ds15MoMsInvT	85%
	5'I3	9Ds16MoBiotin	90%
	5'I4	9Ds16Mp	91%
20	5'I5	9Ds14MoMpD	90%
	5'I6	9Ds13Mo2MpD	94%
	5'I7	9Ds12Mo3MpD	94%
	5'I8	9Ds14MoMsD	93%
	5'I9	9Ds13Mo2MsD	97%
25	5'I10	9Ds12Mo3MsD	95%

Key: M and D refer to 2'O-methyl- and 2'deoxy-ribonucleotides, respectively. The letters "o", "s" and "p" refer to phosphodiester, phosphorothioate diester, and P-ethoxy-phosphotriester linked nucleotides. "InvT" refers to a 3'→3' or 5'→5' linked thymidine at the 3' or 5' end, respectively.

30 Table I shows the results of control oligos C1-C6, all phosphorothioate oligos S1-S4, and oligos of the invention having the RNase activating region at the 3' end (3'I1-3'I3), in the middle (MI1-MI3) and at the 5' end (5'I1-5'I10).

35 Control oligos C1, C2, C5 and C6 showed low levels of specific inhibition because these oligos lacked an RNase H activating region. Oligos C3 and C4 were inactive because

the 3' was unprotected and because native ssDNA was unstable, respectively. All phosphorothioate oligonucleotides (S1-S4) showed specific inhibitions that ranged between 93% and 100%, as did oligonucleotides 5'I6-5'I10, which have a 5'-located 5 RNase H activating region and two or three 3' terminal 2'O-methyl modified P-ethoxy or phosphorothioate linked nucleotides (Mp and Ms, respectively). Lower levels of specific inhibition were observed when oligonucleotides with 3' and mid-located RNase H activating regions were employed 10 or when suboptimal 3' protecting groups were present.

Although the oligonucleotides of the invention having 5' RNase activating regions achieved specific inhibitions which were comparable to that achieved by the uniform phosphorothioate oligonucleotides, the oligonucleotides of 15 the invention were superior in that their use was associated with significantly less toxicity. Table II shows specific inhibition, the average metabolic activity as percent of no oligo control, as determined by MTS assay, and the percent viable cells, as determined by trypan blue exclusion for the 20 conventional ("C"), all phosphorothioate ("S"), 3'I, MI and 5'I oligonucleotides, as well as for three species.

TABLE 2

25	Oligo	% INH Luc	% of Control	% of Viable
			Metabolic Activity	
	All "O" Oligos C1-C6	15%	94%	76%
	All "S" Oligos S1-S4	96%	25%	21%
30	3'I (1-4)	60%	70%	61%
	MI (1-3)	74%	77%	67%
	5'I (1-10)	91%	71%	60%

The best oligos on the chart have high percentage values in 35 all columns.

The results demonstrated that the oligonucleotides of the invention achieve levels of specific inhibition more than

four times greater than conventional oligonucleotides while showing toxicity levels that were substantially less than the phosphorothioate oligonucleotides. The optimal group, 5'I, showed specific inhibition that was comparable to the 5 phosphorothioate oligonucleotides.

5.3 THE EFFECT OF THE LOCATION OF THE RNASE H ACTIVATING REGION

The cause of lower specific activity observed for the 3'I and MI type oligonucleotides was investigated. One possibility was that the oxidation steps using 0.02M I₂ cause the oxidation of the phosphorothioate linkages to phosphodiester, when phosphodiester linked nucleotides were added 5' to the phosphorothioate linkages. This was found to be the case. Comparison of oligonucleotides 9D_s15D_oD ("5'S") and 15D_o9D_sD ("3'S") oligonucleotides having the sequence of the test oligonucleotide by analytical HPLC analysis showed that about 85% of the 5'S oligonucleotides were fully thiolated, by contrast only 26% of the 3'S oligonucleotides were completely thiolated (36% were S-1, 24% S-2 and 14% S-3).

Table III shows the distribution of fully thiolated and mono, di and tri-oxidized by-products as a function of the position of the phosphorothiolated region of the oligonucleotide. Four thymidyl pentadodecamers were synthesized using 0.02 M I₂ as the oxidant for 15 nucleotides and a thiolating agent for nine nucleotides.

TABLE 3

	Ts	[I ₂]	S	S-1	S-2	S-3
30	5'-9Ds15DoD03'	0.02M	96%	4%	-	-
	5'-1Do9Ds14DoD-3'	0.02M	85%	15%	-	-
	5'-8Do9Ds7DoD-3'	0.02M	41%	46%	12.5	0.5
	5'-15Do9DsD-3'	0.02M	32%	43%	20%	5%
35	5'-15Do9DsD-3'	0.001M	78%	14%	8%	-

The results demonstrated that 96% of the 5'S oligonucleotides are fully thiolated, which percentage steadily decreases as the phosphorothioate region is exposed to more frequent oxidation reactions. When the oxidant concentration was reduced to 0.001M, 78% fully thiolated 3'S 25-T oligonucleotides and about 60% of oligonucleotides having the sequence of the SEQ ID NO: 1 were synthesized.

10

15

20

25

30

35

WE CLAIM:

1. A chimeric antisense oligonucleotide comprising: a 5' terminus; a 3' terminus and from 11 to 59 5'→3'-linked nucleotides capable of contiguously hybridizing to a specific RNA and independently selected from the group consisting of 2'-modified phosphodiester nucleotides, and 2'-modified P-alkyloxyphosphotriester nucleotides; and wherein said 11 to 59 5'→3'-linked nucleotides are divided by an RNase H-activating region capable of contiguously hybridizing to the specific RNA and of between three and ten contiguous phosphorothioate-linked deoxyribonucleotides, and wherein the 3' terminus of said oligonucleotide is drawn from the group consisting of: an inverted deoxyribonucleotide, a contiguous stretch of one to three phosphorothioate 2'-modified ribonucleotides, a biotin group, and a P-alkyloxyphosphodiester nucleotide, and wherein the 5' terminus of said oligonucleotide is drawn from the group consisting of: an inverted deoxyribonucleotide, a contiguous stretch of one to three phosphorothioate 2'-modified ribonucleotides, a biotin group, and a P-alkyloxyphosphodiester nucleotide.
2. The oligonucleotide of claim 1, provided the 3' terminus is not blocked by a 3'→3' phosphorothioate linked nucleotide.
3. The oligonucleotide of claim 1, in which the 3' terminus is blocked by a moiety comprising a 3'→3' phosphorothioate linked nucleotide.
4. The oligonucleotide of claim 1, in which the 3' terminus is blocked by a moiety comprising a 3'→3' phosphodiester linked nucleotide.
5. The oligonucleotide of claim 4, in which the 3' most 5'→3' internucleotide linkage is a phosphorothioate linkage or a P-ethoxyphosphotriester linkage.

6. The oligonucleotide of claim 4, in which the 5' most 5'→3' internucleotide linkage is a phosphorothioate linkage or a P-ethoxyphosphotriester linkage.

7. The oligonucleotide of claim 1, in which the 3' terminal nucleoside and the 5' most nucleotide are 2'-modified nucleotides.

8. The oligonucleotide of claim 7, in which the 5' most 5'→3' internucleotide linkage is a phosphorothioate linkage or a P-ethoxyphosphotriester linkage.

9. The oligonucleotide of claim 8, in which the two 5' most 5'→3' internucleotide linkages are independently either a phosphorothioate linkage or a P-ethoxyphosphotriester linkage.

10. The oligonucleotide of claim 8, in which all phosphorothioate linkages are contiguous with the 3' most 5'→3' internucleotide linkage.

11. The oligonucleotide of claim 10, in which the 2'-modified nucleotide is a 2'-methoxy or 2'-fluoro nucleotide.

12. The oligonucleotide of claim 10, which comprises at least thirteen 2'-methoxy phosphodiester nucleotides.

13. The oligonucleotide of claim 10, having between 15 and 50 nucleotides.

14. The oligonucleotide of claim 13, which comprises at least eight 2'-methoxy phosphodiester nucleotides.

15. The oligonucleotide of claim 13, which comprises at least thirteen 2'-methoxy phosphodiester nucleotides.

16. The oligonucleotide of claim 1, in which the 2'-modified nucleotides are selected from the group consisting of 2'-fluoro and 2'-methoxy nucleotides.


17. The oligonucleotide of claim 1, in which there are no 2'-modified phosphorothioate nucleotides.

18. A method of specifically cleaving an RNA in a cell containing RNase H which comprises administering an effective amount of an oligonucleotide complementary to the RNA comprising: a 5' terminus; a 3' terminus; and from 11 to 59 5'→3'-linked nucleotides capable of contiguously hybridizing to the RNA and independently selected from the group consisting of 2'-modified phosphodiester nucleotides, 2'-modified P-alkyloxyphosphotriester nucleotides; and wherein said 11 to 59 5'→3'-linked nucleotides are divided by an RNase H-activating region capable of contiguously hybridizing to the RNA and of between three and ten contiguous phosphorothioate-linked deoxyribonucleotides, and wherein the 3' terminus of said oligonucleotide is drawn from the group consisting of: an inverted deoxyribonucleotide, a contiguous stretch of one to three phosphorothioate deoxyribonucleotides, phosphorothioate 2'-modified ribonucleotides, a biotin group, and a P-alkyloxyphosphodiester-linked nucleotide, and wherein the 5' terminus of said oligonucleotide is drawn from the group consisting of: an inverted deoxyribonucleotide, a contiguous stretch of one to three phosphorothioate deoxyribonucleotides, phosphorothioate 2'-modified ribonucleotides, a biotin group, and a P-alkyloxyphosphodiester-linked nucleotide.

19. A chimeric antisense oligonucleotide comprising:
a) an RNase H activation region capable of contiguously hybridizing to a specific RNA and having between 5 and 10 contiguous deoxyphosphorothioate nucleotides;

- b) between 4 to 59 contiguous 5'→3'-linked 2'-methoxy ribonucleotides capable of contiguously hybridizing to the specific RNA;
- c) an exonuclease blocking group present at the 3' end, the 5' end, or both the 3' and 5' ends of the oligonucleotide drawn from the group consisting of: a non-5'-3' phosphodiester-linked nucleotide, from one to three contiguous 5'-3'-linked modified nucleotides, and a non-nucleotide chemical blocking group.

United States Patent & Trademark Office
Office of Initial Patent Examination -- Scanning Division



Application deficiencies were found during scanning:

☐ Page(s) _____ of DECLARATION were not present
for scanning. (Document title)

☐ Page(s) _____ of _____ were not present
for scanning. (Document title)

☐ Scanned copy is best available.